

BIAŁYCNICKI-BIRULA SCHEMES IN HIGHER DIMENSIONAL HILBERT SCHEMES OF POINTS

LAURENT EVAIN AND MATHIAS LEDERER

ABSTRACT. The Białynicki-Birula cells on the Hilbert scheme $H^n(\mathbb{A}^d)$ are smooth and reduced in dimension $d = 2$. We prove that there is a schematic structure in higher dimension, the Białynicki-Birula scheme, which is natural in the sense that it represents a functor. Let $\rho_i : H^n(\mathbb{A}^d) \rightarrow \text{Sym}^n(\mathbb{A}^1)$ be the Hilbert-Chow morphism of the i^{th} coordinate. We prove that a Białynicki-Birula scheme associated with an action of a torus T is schematically included in the fiber $\rho_i^{-1}(0)$ if the i^{th} weight of T is non positive.

1. INTRODUCTION

Let $H^n(\mathbb{A}^d)$ be the Hilbert scheme parametrizing zero dimensional subschemes of length n in the affine d -space $\mathbb{A}^d$ over a field k . This scheme is mostly called the *Hilbert scheme of points*, sometimes also the *punctual Hilbert scheme*. There is a natural action of the d -dimensional torus on $\mathbb{A}^d$, which induces a natural action on $H^n(\mathbb{A}^d)$. If T is a general one-dimensional subtorus of the d -dimensional torus, then T defines the *Białynicki-Birula schemes* $H^{BB(T,\Delta)}$ parametrizing the subschemes converging to some fixed point Z^Δ under the action of T . The fixed point in question is a monomial subscheme Z^Δ .

These stratifications are preeminent in most studies of the punctual Hilbert scheme in dimension two. For instance, they appear in the computation of the Betti numbers (see [ES87], [ES88]), in the determination of the irreducible components of (multi)graded Hilbert schemes (see [Eva04], [MS10]), or in the study of the ring of symmetric functions via symmetric products of embedded curves (see [Gro96], [Nak99]).

The Białynicki-Birula schemes in $H^n(\mathbb{A}^2)$ are in fact affine cells. In contrast, not much is known on these cells for higher dimensional $\mathbb{A}^d$, and the difficulty to control and describe these cells is probably one of the reasons why the Hilbert scheme of points is still mysterious in higher dimension.

Date: December 2010.

Key words and phrases. Hilbert scheme of points, standard sets, Białynicki-Birula cells after Ellingsrud and Strømme.

The second author was supported by a Marie Curie International Outgoing Fellowship of the EU Seventh Framework Program.

In dimension three or higher, the Białynicki-Birula schemes are not irreducible, nor are they expected to be reduced. It is therefore necessary to define them with their natural scheme structure as representing a functor. Apart from the necessity to define them schematically, it is desirable to have functorial descriptions of Hilbert schemes at hand, as these descriptions are known to be both powerful and easy to handle.

In the present paper, we introduce the Białynicki-Birula functor parametrizing families of subschemes Z such that $\lim_{t \rightarrow 0, t \in T} t.Z = Z^\Delta$ for some fixed monomial subscheme Z^Δ . We will prove (see theorem 25):

Theorem. *The Białynicki-Birula functor is representable by a locally closed scheme $H^{BB(T,\Delta)}(\mathbb{A}^d)$ of the Hilbert scheme $H^n(\mathbb{A}^d)$.*

We assume that the one-dimensional subtorus T of the d -dimensional torus acting on $\mathbb{A}^d$ takes the shape $T = \{(t^{\xi_1}, \dots, t^{\xi_d}), t \in k^*\}$, for suitable weights ξ_i .¹ If $\xi_i \leq 0$, then the closed points of $H^{BB(T,\Delta)}(\mathbb{A}^d)$ correspond to subschemes Z whose support is in the hyperplane $x_i = 0$. This follows from the naïve observation that if $t.Z$ tends to Z^Δ , then the support of Z tends to the support of Z^Δ . A much more subtle question is to ask whether this remains true at the schematic level, when we consider the Białynicki-Birula scheme with its possibly non-reduced structure. The answer is positive. We will prove (see theorem 28):

Theorem. *Let $\rho_i : H^n(\mathbb{A}^d) \rightarrow \text{Sym}^n(\mathbb{A}^1)$ be the Hilbert-Chow morphism which associates to a subscheme Z of length n the unordered n -tuple in k corresponding to the i^{th} coordinate. If $\xi_i \leq 0$, then $H^{BB(T,\Delta)}(\mathbb{A}^d)$ is schematically included in the fiber $\rho_i^{-1}(O)$.*

For simplicity, we have considered a field k in this introduction. But throughout the paper, we shall work over a ring k of arbitrary characteristic.

Let us say a word about the proofs. The action of T on $\mathbb{A}^d = \text{Spec } k[\mathbf{x}]$ induces a partial order $<$ on the monomials of $k[\mathbf{x}]$: the monomials are ordered according to their weight for the T -action. The basic idea of the proof is that a subscheme Z is in $H^{BB(T,\Delta)}$ if, and only if, the initial ideal $\text{in}_{<_\epsilon}(I(Z))$ equals the monomial ideal $I(Z^\Delta)$ for any total order $<_\epsilon$ which is a small deformation of the partial order $<$. It follows that the Białynicki-Birula functor is an intersection of functors $\cap_{<_\epsilon} \mathcal{H}^{\text{mon}(<_\epsilon, \Delta)}$, where the intersection is taken over all small deformations $<_\epsilon$ of $<$, and each functor $\mathcal{H}^{\text{mon}(<_\epsilon, \Delta)}$ is defined using the theory of initial ideals. We prove that each $\mathcal{H}^{\text{mon}(<_\epsilon, \Delta)}$ is representable by a subscheme $H^{\text{mon}(<_\epsilon, \Delta)}$ of $H^n(\mathbb{A}^d)$ (see theorem 21). For avoiding the problems coming from intersecting an infinite number of subvarieties, we realize the Białynicki-Birula functors as the intersection of only two well-chosen functors $\mathcal{H}^{\text{mon}(<_-, \Delta)} \cap \mathcal{H}^{\text{mon}(<_+, \Delta)}$ (see proposition 24).

When dealing with representations of functors, constructions for individual subschemes often require uniformity lemmata when one passes to families. The paragon of this situation is Castelnuovo-Mumford regularity, which appears in the construction of the Hilbert scheme. Our strategy for proving representability of the

¹ The action on closed points of $\mathbb{A}^1$ is thus given by $t.(a_1, \dots, a_d) := (t^{\xi_1} a_1, \dots, t^{\xi_d} a_d)$

Białynicki-Birula functor is no exception; we need two key uniformity lemmata (see lemmata 10 and 14). They are proved and explained in section 3.

Acknowledgements. We give many thanks to Robin Hartshorne, Diane Maclagan and Gregory G. Smith, the organizers of the workshop *Components of Hilbert Schemes* held at American Institute of Mathematics in Palo Alto, CA, in July 2010. It was there that we first met and shared our thoughts; the exceptionally productive and at the same time friendly atmosphere of that workshop triggered our collaboration. Special thanks go to Bernd Sturmfels, who realized that we have many research interests in common, and motivated us to work together. The second author wishes to thank Allen Knutson, Jenna Rajchgot, and Mike Stillman for many fruitful discussions.

The second author was supported by a Marie Curie International Outgoing Fellowship of the EU Seventh Framework Program.

2. BIALYNICKI-BIRULA FUNCTORS AND Δ -MONIC FAMILIES

In this section, we introduce the Białynicki-Birula functor and reformulate Białynicki-Birula families in terms of Δ -monic ideals (see proposition 8).

In this paper, we consider schemes over a commutative ring k of arbitrary characteristic. We denote by $k[\mathbf{x}]$ the polynomial ring $k[x_1, \dots, x_d]$. Similarly, for $e = (e_1, \dots, e_d) \in \mathbb{N}^d$, we use the multi-index notation $\mathbf{x}^e := x_1^{e_1} \dots x_d^{e_d}$. All the rings and k -algebras are implicitly assumed to be noetherian.

A *standard set*, or *staircase*, is a subset $\Delta \subset \mathbb{N}^d$ whose complement $C := \mathbb{N}^d \setminus \Delta$ satisfies $C + \mathbb{N}^d = C$. We call the minimal generators of the $\mathbb{N}^d$ -module C the *outer corners of Δ* . All standard sets under consideration will be of cardinality n , in particular, finite. The ideal generated by the monomials $\mathbf{x}^e, e \in C$ is denoted by I^Δ . We shall freely identify the monomials $\mathbf{x}^e$ with their exponent e . In particular, the notion of a staircase of monomials makes sense.

If B is a k -algebra, then the tensor product $B \otimes_k k[\mathbf{x}]$ is just $B[\mathbf{x}]$, the ring of polynomials with coefficients in B . Similarly, we write $B[t, t^{-1}, \mathbf{x}] := B[\mathbf{x}] \otimes_k k[t, t^{-1}]$. Let $\xi \in \mathbb{Z}^d$, and $I \subset B[\mathbf{x}]$. We denote by $I(t, t^{-1}) \subset B[t, t^{-1}, \mathbf{x}]$ the ideal generated by the elements $t.f := \sum t^{-\xi \cdot e} c_e \mathbf{x}^e$ where $f = \sum c_e \mathbf{x}^e \in I$. We denote by $I(t)$ the ideal $I(t, t^{-1}, \mathbf{x}) \cap B[t, \mathbf{x}]$. In particular $I = I(1)$.

Definition 1. We denote by $\mathcal{H}^{BB(\Delta, \xi)}(B)$, or more simply by $\mathcal{H}^{BB(\Delta)}(B)$ when ξ is obvious, the set of ideals $I \subset B[\mathbf{x}]$ such that $\lim_{t \rightarrow 0} t.I = I^\Delta$, which means:

- $B[t, \mathbf{x}]/I(t)$ is a locally free $B[t]$ -module of rank $n = \#\Delta$.
- $I(0) = I^\Delta$

$\mathcal{H}^{BB(\Delta)}$ is a contravariant functor from the category of noetherian k -algebras to the category of sets. We call it the *Białynicki-Birula functor*.

The first bulleted item of the definition says that $\operatorname{Spec} B[t, \mathbf{x}]/I(t) \rightarrow \operatorname{Spec} B[t]$ is a flat family. The second bulleted item says that its fiber over $0 \in \operatorname{Spec} B[t]$ is the monomial scheme defined by I^Δ . The limit $\lim_{t \rightarrow 0} t.I$ is therefore a well-defined flat limit.

Let us check that $\mathcal{H}^{BB(\Delta, \xi)}$ is indeed a functor. If $Q := B[t, \mathbf{x}]/I(t)$ is a locally free module of rank $\#\Delta$ and $B \rightarrow A$ is a ring homomorphism, then $Q \otimes_B A = Q \otimes_{B[t]} A[t]$ is a locally free $A[t]$ -module of the same rank. In particular, the fiber over $0 \in \operatorname{Spec} A[t]$ of the family $\operatorname{Spec} Q \otimes_B A \rightarrow \operatorname{Spec} A[t]$ is a locally free A -module, call it Q' , of the same rank. However, the kernel of the corresponding surjection $A[\mathbf{x}] \rightarrow Q'$ contains the monomial ideal I^Δ . Since the quotient $A[\mathbf{x}]/I^\Delta$ is a free A -module of rank $\#\Delta$, the module Q' is in fact free of that rank; the kernel of $A[\mathbf{x}] \rightarrow Q'$ equals I^Δ ; and functoriality follows.

We shall now introduce the partial order on monomials in $k[\mathbf{x}]$ defined by the weight of our torus action. That order is in general not a total order, in particular, not a monomial order. In what follows, we shall employ techniques very similar to those in Gröbner basis theory—with the difference that our order is not a monomial order, so the usual techniques will have to be modified. Let us make that more precise.

Definition 2. Let $\xi \in \mathbb{R}^d$ and $f_\xi \in (\mathbb{R}^d)^*$ the linear form defined by $f_\xi(\alpha_1, \dots, \alpha_d) = \sum \alpha_i \xi_i$. We define the partial order $<_\xi$ on monomials in $k[\mathbf{x}]$ by setting $\mathbf{x}^e <_\xi \mathbf{x}^g$ if $f_\xi(e) < f_\xi(g)$, and letting $\mathbf{x}^e$ and $\mathbf{x}^g$ be incomparable if $f_\xi(e) = f_\xi(g)$. Since we identify monomials and exponents, we adopt the convention $f_\xi(x^e) := f_\xi(e)$.

If the weights ξ_i are linearly independent over $\mathbb{Q}$, then $<_\xi$ is a total order on monomials. Otherwise, the order is only partial. In that case we shall refine the partial order to a total one. The refinements we shall use may be defined either as limits or using *signed orders*.

Definition 3.

- A sequence of partial orders $<_j$ *converges to the total order* $<$ if for every pair of monomials a, b , we have $a < b$ if, and only if, $a <_j b$ for j large enough.
- A *signed order* on the symbols $1, \dots, d$ is a map $(\epsilon, o) : \{1, \dots, d\} \rightarrow \{-1, 1\} \times \{1, \dots, d\}$ such that the induced map $o : \{1, \dots, d\} \rightarrow \{1, \dots, d\}$ is a bijection.
- A *sequence compatible with that signed order* is a sequence ξ^j in $\mathbb{R}^d$ converging to 0 such that the sign of ξ_k^j is $\epsilon(k)$ and such that the quotient ξ_i^j / ξ_k^j tends to 0 if, and only if, $o^{-1}(k) < o^{-1}(i)$.

The connection between signed orders and convergence is as follows.

Proposition 4. *Let $<$ be a refinement of the order $<_\xi$ and (ϵ, o) a signed order. Then the following conditions are equivalent:*

- For all monomials $\mathbf{x}^e$ and $\mathbf{x}^f$ with $f_\xi(e) = f_\xi(f)$, we have $\mathbf{x}^e < \mathbf{x}^f$ if, and only if, $(\epsilon(1)e_{o(1)}, \dots, \epsilon(d)e_{o(d)}) < (\epsilon(1)f_{o(1)}, \dots, \epsilon(d)f_{o(d)})$ in the lexicographic order.
- For every sequence ξ^j compatible with the signed order, the sequence of orders $<_{\xi+\xi^j}$ converges to $<$.

Proof. Easy. $\square$

Definition 5. A *quasi-homogeneous order of weight ξ* is either a partial order $<_\xi$ as above or a total order $<$ which refines $<_\xi$. The variable x_i is *positive* with respect to $<_\xi$ if $\xi_i > 0$. If the total order $<$ refines $<_\xi$, then a variable x_i with $\xi_i = 0$ and $\epsilon(i) = 1$ in the refinement is called *positive* too. This defines positivity of variables for all quasi-homogeneous order of weight ξ . *Negativity* of variables is defined similarly.

When all variables are positive with respect to a total order $<$, then the order is a monomial order and the theory of Gröbner bases is at our disposition. In particular, there exist an algorithm for division of an element f by a monic family f_i . When all variables are negative, we are in the setting of standard bases, in which there also exists a division algorithm. With standard bases, the quotient is a power series. However, the quotient is a polynomial when the family f_i contains a power x_j^k of each variable x_j . In what follows we shall consider weight vectors ξ whose signs of coordinates are neither all positive nor all negative. In this setting, there exists a division algorithm with polynomial quotient by a monic family f_i containing powers of the negative variables. This leads to the notion of a *bounded ideal*.

Definition 6. Let $<$ be a quasi-homogeneous total order. An ideal $I \subset B[\mathbf{x}]$ is called *bounded by $r_1, \dots, r_d$* if for every negative (resp. positive) variable x_i , there exists a polynomial $h_i \in I \cap B[x_i]$ with $h_i = x_i^{r_i}$ (resp. $\text{in}(h_i) = x_i^{r_i}$).

Definition 7. • Let $<$ be a quasi-homogeneous partial order of weight ξ , and $f = \sum a_e \mathbf{x}^e \in B[\mathbf{x}]$, where $a_e \neq 0$ for every e . Let $\mathbf{x}^{e_1}, \dots, \mathbf{x}^{e_l}$ be the maximal monomials appearing in f . The *initial form of f* is $\text{in}(f) := \sum a_{e_i} \mathbf{x}^{e_i}$. This is a term when $<$ is a total order, but may contain more than one term otherwise. We denote by $\text{in}(I)$ the ideal generated by the elements $\text{in}(f)$, $f \in I$.

- Let $I \subset B[\mathbf{x}]$ and $m = \mathbf{x}^e$ be a monomial. We denote by $\text{in}_m(I) \subset B$ the ideal generated by the elements $b \in B$ such that $\text{in}(f) = bm$ for some $f \in I$.
- Let Δ be a standard set of cardinality n . The ideal I is called Δ -monic if $\text{in}_m(I) = \langle 1 \rangle$ if $m \notin \Delta$ and $\text{in}_m(I) = 0$ otherwise.

We now interpret the Białynicki-Birula functor in terms of initial ideals and monic families with respect to some partial quasi-homogeneous order.

Proposition 8. Let $\xi \in \mathbb{Z}^d$ and $<_\xi$ the associated partial order. Then $I \in \mathcal{H}^{BB(\Delta, \xi)}(B)$ if, and only if, the following conditions are satisfied:

- I is Δ -monic
- $\forall f \in I$, the initial form with respect to $<_\xi$ is $\text{in}(f) = \sum a_e x^e$, $e \notin \Delta$,
- $B[\mathbf{x}]/I$ is a locally free B -module of rank $\#\Delta$.

Proof. We only prove that the itemized conditions imply that $I \in \mathcal{H}^{BB(\Delta, \xi)}$, the converse being easy. Let us temporarily assume that $B[t, \mathbf{x}]/I(t)$ is a locally free $B[t]$ -module. Then the flat limit $\lim_{t \rightarrow 0} t \cdot I$ exists; the first condition says that the limit ideal contains I^Δ , whereas the second conditions says that the limit is

contained in I^Δ . For proving that $I \in \mathcal{H}^{BB(\Delta, \xi)}(B)$, it therefore remains to show that $P := B[t, \mathbf{x}]/I(t)$ is a locally free $B[t]$ -module.

Upon localizing B (that is to say, upon replacing $\text{Spec } B$ by an open subset), we may assume that $B[\mathbf{x}]/I$ is B -free of rank $\#\Delta$. According to the second item, the monomials $\mathbf{x}^e, e \in \Delta$ are a basis of $B[\mathbf{x}]/I$. These monomials are therefore also a basis of the $B[t, t^{-1}]$ -module $B[\mathbf{x}]/I \otimes_k k[t, t^{-1}] \simeq B[\mathbf{x}, t, t^{-1}]/I(t, t^{-1})$, the latter isomorphism being given by the torus action $t \cdot \mathbf{x}^e = t^{-\xi \cdot e} \mathbf{x}^e$ on the polynomial ring², which induces the torus action on $H^n(\mathbb{A}^d)$. The monomials $\mathbf{x}^e$, for $e \in \Delta$, remain linearly independent in the $B[t]$ -module P . We denote by $N := \langle \mathbf{x}^e : e \in \Delta \rangle$ the free submodule of $B[t, \mathbf{x}]/I(t)$ generated by monomials with exponents in Δ , and by $Q(t)$ the quotient $P/N = B[t, \mathbf{x}]/(I(t) + \langle \mathbf{x}^e : e \in \Delta \rangle)$. Since $I(0) = I^\Delta$ by hypothesis, we have $Q(0) = Q(t)/\langle t \rangle = 0$. By the above, we have $Q(t, t^{-1}) = Q(t) \otimes_{B[t]} B[t, t^{-1}] = 0$. Hence $Q(t) = 0$, so $P = N$. $\square$

Remark 9. *According to the intuition from Gröbner bases, one could think that the third condition is a consequence of the first two. This is not the case, as is shown by the example $d = 1$, $I = \langle x^2 + x^3 \rangle$, $\xi = -1$, $\Delta = \{1, x\}$.*

3. UNIFORMITY LEMMATA

The goal of this section is to prove the uniformity lemmata needed in our constructions.

Uniformity considerations are a cornerstone of many constructions with representable functors. For instance, in the construction of the Hilbert scheme, a subscheme with Hilbert polynomial P is characterized by its equations in degree m for m large enough. But finding a uniform integer m suitable for all the subschemes with Hilbert polynomial P requires a uniformity lemma. In the construction of Hilbert scheme, Castelnuovo-Mumford regularity, or the Gotzmann persistence theorem does the job. In our context, we shall need two uniformity lemmata (see lemmata 10 and 14). Let us explain where they are needed.

In proposition 8, we have reformulated the property $I \in \mathcal{H}^{BB(\xi, \Delta)}(B)$ in terms of the partial order $<_\xi$ and of the ideal $\text{in}_{<_\xi}(I)$. Later on, we shall use total orders instead of partial orders to connect the Białynicki-Birula functors with Gröbner basis theory. We shall slightly deform ξ into ξ_ϵ in such a way that $<_{\xi_\epsilon}$ is a total order and $\text{in}_{<_\xi}(I) = \text{in}_{<_{\xi_\epsilon}}(I)$. But when $<_j$ converges to $<$, in the equivalence $a < b \Leftrightarrow a <_j b$ for j large, how large j needs to be depends on the monomials a and b . The first uniformity lemma (lemma 10) controls the monomials appearing in a nice set of generators of I and will induce the equality $\text{in}_{<_\xi}(I) = \text{in}_{<_{\xi_\epsilon}}(I)$ (corollary 11).

As for the second uniformity lemma (see lemma 14), the situation is as follows. If $Z_0 \subset \mathbb{A}^d$ is a scheme of length n supported on the locus $x_i = 0$, then x_i^n vanishes

² Remember that the torus takes the shape $T = \{(t^{\xi_1}, \dots, t^{\xi_d}), t \in k^*\}$ and acts by $t \cdot (a_1, \dots, a_d) := (t^{\xi_1} a_1, \dots, t^{\xi_d} a_d)$ on closed points of $\mathbb{A}^d$. This corresponds to the torus action on the polynomial ring $k[\mathbf{x}]$ which is trivial on scalars and is given by $t \cdot \mathbf{x}^e := t^{-\xi \cdot e} \mathbf{x}^e$ on monomials.

on Z . This is not true any more in families. The simple example $Z = V(x - \epsilon) \subset \text{Spec } k[x, \epsilon]/\epsilon^2$ shows a family of relative length 1, supported on $x = 0$, schematically included in $x^2 = 0$ but not schematically included in $x = 0$. This corresponds to the fact that the functor parametrizing families of length n supported on $x_i = 0$ is not representable by a closed subscheme of $H^n(\mathbb{A}^d)$. In contrast, we shall prove in lemma 14 that a Δ -monic bounded family of relative length n supported on $x_i = 0$ is schematically included in the locus $x_i^n = 0$. This is an ingredient to prove that the monic functors are representable by locally closed subschemes of $H^n(\mathbb{A}^d)$ (see theorem 21).

Lemma 10. *Let $I = \langle f_1, \dots, f_u \rangle \subset B[\mathbf{x}]$ be an ideal such that the quotient $B[\mathbf{x}]/I$ is a finite B -module. Let $<$ be a quasi-homogeneous total order of weight $\xi = (\xi_1, \dots, \xi_d)$ with $\sum \xi_i^2 = 1$. Assume moreover that one of the following two conditions is satisfied:*

- (1) $\xi_i \neq 0$ for all i ,
- (2) I is bounded.

Then there exists $g_i \in I$ such that for all m and for every $a \in \text{in}_m(I)$, there exist $\lambda_i \in B[\mathbf{x}]$, $b, c \in \mathbb{R}^+$ such that

- $\text{in}(\sum \lambda_i g_i) = am$,
- *the total degree of each term τ in $\sum \lambda_i g_i$ satisfies*
 - case (1): $\deg(\tau) \leq \frac{b}{\min(|\xi_j|)} + c$,
 - case (2): $\deg(\tau) \leq b$,
- b, c depend on I and m , but not on the order $<$.

Proof. The sequence $\langle 1, x_i, \dots, x_i^j \rangle \subset B[\mathbf{x}]/I$, indexed by j , is an increasing sequence in the noetherian module $B[\mathbf{x}]/I$, thus it eventually becomes stationary. It follows that for every i , the ideal I contains an element $h_i = x_i^r + \sum_{j=1}^{r-1} b_{ij} x_i^j$, $b_{ij} \in B$. We define

$$g_i := \begin{cases} f_i & \text{for } i \leq u, \\ h_{i-u} & \text{for } i = u+1, \dots, u+d. \end{cases}$$

Consider an element f of I , which takes the shape

$$f = \sum s_i f_i,$$

and let $\text{in}(f) = am$, where m is a monomial and $a \in B$.

Let $s_i = h_1 q_i + r_i$ be the euclidean division of s_i by h_1 with respect to the variable x_1 . Then

$$f = \sum r_i f_i + (\sum q_i f_i) h_1,$$

where $\deg_{x_1} r_i < r$ for some natural number r which is independent of f . We repeat euclidean division for all other h_j , so with respect to all other variables x_j . This yields a formula

$$f = \sum t_i f_i + \sum u_i h_i,$$

where

$$\deg_{x_j} t_i < r$$

for all i, j and a uniform bound r . We define $D := \max_{i,j} \{\deg_{x_j} f_i + r - 1, \deg_{x_j} m\}$. Let s be the index of a positive variable. Then $\deg_{x_s} \sum t_i f_i \leq D$. Let τ be a term of $\sum u_i h_i$ with $\deg_{x_s} \tau > D \geq \deg_{x_s}(m)$. Then τ is also a term of f , different from its initial term, $\tau \neq \text{in}(f) = am$. Let $f' := f - \frac{\tau}{x_s^r} h_s$. Then $\text{in}(f') = \text{in}(f)$ and

$$f' = \sum t_i f_i + \sum u'_i h_i, u'_s = u_s - \frac{\tau}{x_s^r} h_s, u'_i = u_i, i \neq s.$$

The term τ , which appears in $\sum u_i h_i$, is gone in $\sum u'_i h_i$. We may repeat this construction, thus constructing expressions $\sum u_i^{(l)} h_i$ which suppress all the terms whose degree in some variable of positive weight is more than D . Finally, we obtain an element $f^{(l)} \in I$ such that $\text{in}(f^{(l)}) = am$ and

$$f^{(l)} = \sum t_i f_i + \sum v_i h_i, \quad \deg_{x_j} \sum v_i h_i \leq D \text{ when } \xi_j > 0.$$

Now let s be an index with $\xi_s < 0$. In particular, x_s is a negative variable. Let $\tau := cx^e$ be a term of $\sum v_i h_i$ with $\deg_{x_s} \tau > \frac{f_\xi(m) - dD}{\xi_s} + r$. In particular, $\deg_{x_s} \tau > r$ since $f_\xi(m) \leq \sum_{\xi_i > 0} \xi_i \deg_{x_i} m \leq \sum_{\xi_i > 0} D \leq dD$, and the denominator ξ_s is negative. Let $q_s := \frac{\tau}{x_s^r}$. Note that $f_\xi(\text{in}(q_s h_s)) \leq f_\xi(\text{in}(q_s)) = f_\xi(\tau) - r\xi_s < \xi_s \deg_{x_s} \tau + \sum_{\xi_i > 0} \xi_i \deg_{x_i} \tau - r\xi_s \leq \frac{f_\xi(m) - dD}{\xi_s} \xi_s + \sum_{\xi_i > 0} D < \frac{f_\xi(m) - dD}{\xi_s} \xi_s + dD = f_\xi(m)$. In particular, $\text{in}(f^{(l)}) = \text{in}(f^{(l)} - h_s q_s) = am$. We therefore let $f^{(l+1)} := f^{(l)} - h_s q_s$ and obtain have the formula

$$f^{(l+1)} = \sum t_i f_i + \sum w_i h_i,$$

which has the virtue that the term τ does not appear any more in $\sum w_i h_i$. Repeating the same process for all s such that $\xi_s < 0$, we obtain an element

$$f^{(l+p)} = \sum t_i f_i + \sum z_i h_i$$

such that any term τ of $\sum z_i h_i$ satisfies the inequality

$$\deg_{x_s} \tau \leq \frac{f_\xi(m) - dD}{\xi_s} + r \leq \frac{\sum \deg_{x_i}(m) + dD}{|\xi_s|} + r$$

for every s with $\xi_s < 0$.

Summing things up, in the last expression, any term τ of the right hand side satisfies

- $\deg_{x_i} \tau \leq b_i$ for some b_i if $\xi_i > 0$.
- $\deg_{x_i} \tau \leq \frac{b_i}{|\xi_i|} + c_i$ for some b_i, c_i if $\xi_i < 0$.

Thus we have the required expression in case (1).

It remains to cover the case when I is bounded. In the above proof, the upper bound for $\deg_{x_i}(\tau)$ took two different forms, depending on the sign of ξ_i . This is necessary because the sign of ξ_i affects the computation of $\text{in}(h_i)$. However, in the bounded case, one can take $h_i := x_i^{r_i}$ for some r_i if $\xi_i \leq 0$ is a negative variable, and obtains $\text{in}(h_i) = x_i^{r_i}$. In other words, in the bounded case, with this choice of

h_i , the above proof for positive variables also works for negative variables. We get the upper bound $\deg_{x_i} \tau \leq b_i$ whatever the sign of ξ_i is. $\square$

Corollary 11. *Let Δ be a staircase. Let $<_j$ be a sequence of quasi-homogeneous total orders converging to $<$ and $I \subset B[\mathbf{x}]$ be an ideal such that $B[\mathbf{x}]/I$ is a finite B -module and $\text{in}_m(I) = \langle 1 \rangle$ for $m \notin \Delta$. Assume that one of the following two conditions hold:*

- (1) $\xi_i \neq 0$ for all i ,
- (2) I is bounded.

Then for large j , we have $\text{in}_<(I) = \text{in}_{<_j}(I)$.

Proof. If $\text{in}_<(f) = am$, then $\text{in}_{<_j} f = am$ for large j by definition of the limit order. The inclusions $\text{in}_{m,<}(I) \subset \text{in}_{m,<_j}(I)$ and $\text{in}_<(I) \subset \text{in}_{<_j}(I)$ for large j then follow from the noetherian property.

Conversely, let ξ_j and ξ the normalized weights associated to $<_j$ and $<$, where normalization means that $\|\xi_j\|_2 = \|\xi\|_2 = 1$. Since $<_j$ converges to $<$, the sequence ξ_j converges to ξ . By the previous proposition, for all m , there exists $N_m \in \mathbb{N}$ and a finite family of monomials m_i such that for $j > N_m$, any $a \in \text{in}_{m,<_j}(I)$ is equal to $\text{in}_{<_j}(f)$ for some $f = \sum b_i m_i$, $b_i \in B$. Let $N'_m \in \mathbb{N}$ be such that for all $n > N'_m$, the partial order $<_j$ and the total order $<$ define the same total order on the finite set of monomials m_i . Then for $n > N''_m := \max(N, N')$, we have $\text{in}_{m,<_j}(I) \subset \text{in}_{m,<}(I)$.

When trying to find an integer large enough so that $\text{in}_<(I) = \text{in}_{<_j}(I)$, one a priori thinks that this integer should be at least equal to N_m for every monomial $m \in k[\mathbf{x}]$. But if $\text{in}_{m,<_j}(I) = \text{in}_{m,<}(I) = \langle 1 \rangle$, then the same equality is also still true if one replaces m by any monomial multiple m' of m . In particular, it suffices to take $j > N := \max\{N_m : m \in A\}$, where A is the union of Δ and the outside corners of Δ , a finite set. $\square$

We now provide a (pseudo-)division algorithm which, in particular, tests ideal membership. Of course such an algorithm exists in the presence of a monomial order or with bounded families—but since in our context we need partial and total non monomial orders, that algorithm is not good enough for us. We deform the partial order into a total order and start a division with the total order. Our algorithm does not necessarily terminate. Nevertheless we are able to identify a part R_Δ in the remainder which stabilizes (and is independent of the arbitrarily chosen deformed order). This division is used to prove that most monic families are bounded (corollary 13). Here is the precise statement.

Corollary 12. *Let $<$ be a quasi-homogeneous partial order of weight ξ , Δ a staircase, and $I \subset B[\mathbf{x}]$ a Δ -monic ideal such that the quotient $B[\mathbf{x}]/I$ is locally free of rank $\#\Delta$. Let $o_1, \dots, o_u$ the outside corners of Δ , and $f_1, \dots, f_u \in I$ be elements with $\text{in}_<(f_i) = o_i$. Let ξ' a small deformation of ξ such that $<_{\xi'}$ is a total order and $\text{in}_<(I) = \text{in}_{<'}(I)$ and $\text{in}_<(f_i) = \text{in}_{<'}(f_i)$ for all i . Then for all $f \in B[\mathbf{x}]$, there exists a division*

$$f = \sum \lambda_i f_i + R_\Delta + R'$$

such that

- each term $\tau = b\mathbf{x}^e$ of R_Δ satisfies $e \in \Delta$,
- for all terms $c\mathbf{x}^s$ of R' and for all $m \in \Delta$, $f_{\xi'}(s) < f_{\xi'}(m)$.

For every such division,

- R_Δ is independent of ξ' and of the choice of the division,
- $f \in I$ if, and only if, $R_\Delta = 0$,
- the map $f \mapsto R_\Delta$ is a homomorphism of B -modules $B[\mathbf{x}] \rightarrow B[\mathbf{x}]/I$, where we identify the latter module with $B[\Delta] := \bigoplus_{e \in \Delta} B\mathbf{x}^e$.

Proof. To construct the expected expression $f = \sum \lambda_i f_i + R_\Delta + R'$, we proceed in several steps. At each step j , we have an expression $f = \sum \lambda_{ij} f_i + T_j$. For $j = 0$, we take $\lambda_{i0} := 0$, $T_0 := f$. We decompose $T_j = R_{\Delta,j} + R'_j$, with $R_{\Delta,j} := \sum_{m \in \Delta} c_{mj} m$ and $R'_j := \sum_{m \notin \Delta} c_{mj} m$. The initial term $\text{in}_{<\xi'}(R'_j)$ can be written $\mu_j \text{in}_{<\xi'}(f_{i_j})$ for some f_{i_j} by hypothesis. We set $\lambda_{i_j(j+1)} := \lambda_{i_j j} + \mu_j$ and $\lambda_{i'(j+1)} := \lambda_{i' j}$ for $i' \neq i_j$. Then $T_{j+1} := f - \sum \lambda_{i(j+1)} f_i = T_j - \mu_j f_{i_j}$ decomposes, analogously as above, into $T_{j+1} = R_{\Delta,j+1} + R'_{j+1}$.

If, for some j , it happens that $R'_j = 0$, then we define $R_\Delta := R_{\Delta,j}$ and $R' := 0$, and have constructed the expected expression. Otherwise, there is a constant $c > 0$ such that for all j , $f_{\xi'}(\text{in}_{<\xi'}(f_{i_j})) > f_{\xi'}(\text{in}_{<\xi'}(f_{i_j} - \text{in}_{<\xi'} f_{i_j})) + c$. If we let $E_j := \{f_{\xi'}(\tau) : \tau \text{ a term of } R'_j\}$, then E_{j+1} is derived from E_j by the replacement of the maximal element $M_j = f_{\xi'}(\text{in}_{<\xi'}(R'_j))$ of E_j by a collection $\{c_1, \dots, c_k\}$ with $c_i < M_j - c$ for all i . It follows that for large j , the maximal element of E_j is smaller than any fixed number. In particular, $\forall m \in \Delta$, $M_j = f_{\xi'}(\text{in}_{<\xi'}(R'_j)) < f_{\xi'}(m)$ if j is large enough.

If $f \in I$, then in the expression $f = \sum \lambda_i f_i + R_\Delta + R'$, we have $R_\Delta = 0$ since $\text{in}_{<\xi'}(f - \sum \lambda_i f_i)$ cannot lie in Δ by hypothesis. In particular, if $f = \sum \mu_i f_i + S_\Delta + S'$ is another division, we take their difference and obtain a division of $0 \in I$, which implies $R_\Delta = S_\Delta$.

It is obvious that $f \mapsto R_\Delta$ is a homomorphism as it is possible to add divisions, or to multiply them with a scalar $\lambda \in B$.

Let us now consider the B -submodule $B[\Delta]$ of $B[\mathbf{x}]$. The identity on $B[\Delta]$ factors as $B[\Delta] \rightarrow B[\mathbf{x}] \rightarrow B[\Delta]$ where the first arrow is the inclusion and the second is the morphism R_Δ . The above implies that this factorization induces a factorization $B[\Delta] \rightarrow B[\mathbf{x}]/I \rightarrow B[\Delta]$. This composition is surjective between locally free modules of the same rank, so it is an isomorphism. In particular, we obtain that $R_\Delta = 0$ implies $f \in I$.

To prove that in the division $f = \sum \lambda_i f_i + R_\Delta + R'$, the summand R_Δ is independent of ξ' , it suffices to consider two small deformations ξ' and ξ'' of ξ and to find a division which is valid for both ξ' and ξ'' . We therefore have to show that for all exponents s of terms of R' and for all $m \in \Delta$, $f_{\xi'}(s) < f_{\xi'}(m)$ and $f_{\xi''}(s) < f_{\xi''}(m)$.

Choose a division $f = \sum \lambda_i'' f_i + R_\Delta'' + R''$ which is valid for ξ'' . Then we construct the division for ξ' as above, except for one modification: For $j = 0$, instead of taking $\lambda_{i0} = 0$, $T_0 = f$, we set $\lambda_{i0} := \lambda_i''$, $T_0 := R_\Delta'' + R''$. The terms in R_j' decrease for both $<_{\xi'}$ and $<_{\xi''}$, thus at each step, the expression $f = \sum \lambda_{in} f_{ij} + R_{\Delta,j} + R_j$ is a division for ξ'' . When the process stops, we have a common division for ξ' and ξ'' . $\square$

Corollary 13. *Let ξ a weight vector with $\xi_i \neq 0$ for all i . If I is Δ -monic and $B[\mathbf{x}]/I$ is locally free of rank $n = \#\Delta$, then I is bounded.*

Proof. If r is large and x_i is a negative variable, the division of x_i^r by the collection of f_i reads

$$x_i^r = \sum 0 \cdot f_i + (R_\Delta = 0) + (R' = x_i^r).$$

Thus $x_i^r \in I$, as expected. If x_i is a positive variable, the polynomial h_i constructed at the beginning of lemma 10 is the required polynomial. $\square$

We now come to the second uniformity lemma.

Lemma 14. *Let $<$ be a quasi-homogeneous total order. Let $I \subset B[\mathbf{x}]$ be a Δ -monic ideal with $B[\mathbf{x}]/I$ locally free of rank $n = \#\Delta$. Suppose that I is bounded. Then for every negative variable x_i , we have $x_i^n \in I$.*

Proof. After applying a suitable permutation, we may assume that $x_1, \dots, x_l$ (resp. $x_{l+1}, \dots, x_d$) are the negative (resp. positive) variables, and we shall prove that $x_1^n \in I$. For every monomial $m \in k[x_2, \dots, x_d]$, there exists a unique $h(m) \in \mathbb{N}$, call it the *height*, such that $m^- := x_1^{h(m)-1} m \in \Delta$ and $m^+ := x_1^{h(m)} m \notin \Delta$. In particular, $h(m) = 0$ if $m \notin \Delta$, and all heights of all $\mathbf{x}^e \in k[x_2, \dots, x_d]$ sum up to n , the cardinality of Δ .

Let $M \subset k[x_2, \dots, x_l]$ be the set of monomials with exponents in $\Delta' := \mathbb{N}^{l-1} \cap \Delta$. The set M is finite, and we number its elements such that $m_1^+ < m_2^+ < \dots < m_{\#M}^+$. For any monomial $m \in k[x_2, \dots, x_d]$, we define $H(m) := \sum_{m_i^+ \leq m^+} h(m_i)$. Note that for every such m , $H(m) \leq \sum_{\mathbf{x}^e \in k[x_2, \dots, x_d]} h(\mathbf{x}^e) = n$. In particular, if we prove

$$(1) \quad \forall m = \mathbf{x}^e \in k[x_2, \dots, x_l], x_1^{H(m)} m \in I$$

then for the particular choice $m := 1$, (1) implies $x_1^n \in I$, which concludes the proof.

Since I is bounded, for each i with $2 \leq i \leq l$, there exists some r_i such that $x_i^{r_i} \in I$. Thus (1) is true if $x_i^{r_i}$ divides m . It follows that the set $C := \{m \in k[x_2, \dots, x_l] : m \text{ is a monomial, } x_1^{H(m)} m \notin I\}$ is finite. If (1) is not true, then $C \neq \emptyset$ and C contains an element m_0 which is minimal in the sense that $m_0^+ < m^+$ for every $m \in C, m \neq m_0$. Let $f \in I$ with $\text{in}(f) = m_0^+$. Then f takes the shape

$$(2) \quad f = m_0^+ + \sum_{\mathbf{x}^e < m_0^+} c_e \mathbf{x}^e.$$

Since I is bounded, we may assume by the standard division procedure that $\mathbf{x}^e \in \Delta$ for every term $\mathbf{x}^e$ in the above expression. We obtain

$$\mathbf{x}^e = x_1^{e_1} \dots x_d^{e_d} \in \Delta \Rightarrow x_1^{e_1} \dots x_l^{e_l} \in \Delta \Rightarrow e_1 < h(x_2^{e_2} \dots x_l^{e_l}).$$

Since x_1 is a negative variable, and $x_{l+1}, \dots, x_d$ are positive, we get:

$$(x_2^{e_2} \dots x_l^{e_l})^+ = x_1^{h(x_2^{e_2} \dots x_l^{e_l})} (x_2^{e_2} \dots x_l^{e_l}) < x_1^{e_1} x_2^{e_2} \dots x_l^{e_l} x_{l+1}^{e_{l+1}} \dots x_d^{e_d} = \mathbf{x}^e$$

Here is the upshot of the above: If $C \neq \emptyset$, $m_0 \in C$ is its minimum, and a monomial $\mathbf{x}^e = x_1^{e_1} \dots x_d^{e_d}$ appears in a term of f , then $m_e := x_2^{e_2} \dots x_l^{e_l}$ satisfies $m_e^+ < \mathbf{x}^e < m_0^+$. Thus $H(m_e) \leq H(m_0) - h(m_0)$, and by minimality of m_0 , $m_e x_1^{H(m_e)} \in I$. It follows that the multiple $\mathbf{x}^e x_1^{H(m_0) - h(m_0)}$ lies in I . The product of the expression (2) with $x_1^{H(m_0) - h(m_0)}$ yields

$$m_0 x_1^{H(m_0)} = m_0^+ x_1^{H(m_0) - h(m_0)} = f x_1^{H(m_0) - h(m_0)} - \sum c_e \mathbf{x}^e x_1^{H(m_0) - h(m_0)} \in I,$$

a contradiction. It follows that $C = \emptyset$, and (1) is true. $\square$

4. BASE CHANGE FOR MONIC FAMILIES

It is well-known that passage to the initial ideal does not commute with arbitrary base change, but rather only with flat base change (see [BGS93]). Thus it is difficult to define functors manipulating initial ideals through arbitrary base changes. In contrast, we show that Δ -monic families are stable by arbitrary base change. This allows us to define a Δ -monic functor at the end of this section, after having settled the required base-change statements.

Proposition 15. *Let $<$ be a quasi-homogeneous total order. Let $I \subset B[\mathbf{x}]$ be a bounded ideal. If for all m , $\text{in}_m(I) = 0$ or $\text{in}_m(I) = \langle 1 \rangle$, then for every base change $B \rightarrow A$, $\text{in}_m(IA[\mathbf{x}]) = \text{in}_m(I)A$.*

Proof. Obviously, $\text{in}_m(I) = \langle 1 \rangle$ implies $\text{in}_m(IA[\mathbf{x}]) = \langle 1 \rangle$. If $\text{in}_m(I) = 0$, we argue by contradiction, supposing that $\text{in}_m(IA[\mathbf{x}]) \neq 0$. Choose $f \in IA[\mathbf{x}]$ with $\text{in}(f) = am$, $a \neq 0$. Let x_j be a negative variable. Then $x_j^{d_j} \in I$ for d_j large. In particular, replacing f with $f - \lambda x_j^e$, we may assume that f has no term divisible by $x_j^{d_j}$. Choose an expression

$$(3) \quad f = \sum a_i f_i$$

with $a_i \in A$ and $f_i \in I \subset B[\mathbf{x}]$. As above, we may assume that none of the f_i contains a term divisible by $x_j^{d_j}$. Let m' be the maximal monomial appearing in the f_i . Note that $m' \geq m$ and, more precisely, $m' > m$ since $\text{in}_m(I) = 0$. Among all possible expressions (3), choose one with minimal m' . (Although $<$ is not necessarily a monomial order, a minimum m' exists since we have bounded the exponent of the negative variables). Let $J := \{i : \text{in}(f_i) = \lambda_i m'\}$ be the set of indices of f_i with initial monomial m' . The coefficient of m' vanishes on the right hand side of (3), thus $\sum_{i \in J} a_i \text{in}(f_i) = \sum_{i \in J} (a_i \lambda_i) m' = 0$. Thus $\sum_{i \in J} a_i \lambda_i = 0$.

Let $g \in I$ with $\text{in}(g) = m'$ and such that g has no term divisible by $x_j^{d_j}$. Let $f'_i := f_i$ if $i \notin J$ and $f'_i := f_i - \lambda_i g$ if $i \in J$. Then

$$f = \sum a_i f'_i$$

and this expression contradicts the maximality of m' . $\square$

We remark that with the same proof, we have in fact a slightly more precise and technical statement which will be useful later.

Proposition 16. *Let $<$ be a quasi-homogeneous total order. Let $I \subset B[\mathbf{x}]$ be a bounded ideal. Let m be a monomial. If for all $m' \geq m$, either $\text{in}_{m'}(I) = 0$ or $\text{in}_{m'}(I) = \langle 1 \rangle$ holds, then for every base change $B \rightarrow A$, the equality $\text{in}_m(\text{in}_m(I)A) = \text{in}_m(I)A$ holds.*

Let $X = \text{Spec } A$ be the affine scheme corresponding to the ring A . The ideal $\text{in}_m(I) \subset A$ defines a closed subscheme of X . If X is a scheme which is not affine, we wish to glue the local constructions we have been working with so far. Since open immersions are flat, the following proposition implies that gluing is possible and that for any sheaf of ideals $\mathcal{I} \subset \mathcal{O}_X[x]$, there is a well-defined sheaf of ideals $\text{in}_m(\mathcal{I}) \subset \mathcal{O}_X$ on a non-affine scheme X . This allows us to speak of *bounded* and *monic*, resp., *ideal sheaves* rather than ideals.

Proposition 17. *Let $<$ be a quasi-homogeneous total order. Let $I \subset B[\mathbf{x}]$ be a bounded ideal. Let $B \rightarrow A$ be a ring homomorphism which makes A a flat B -module. Then $\text{in}_m(\text{in}_m(I)A) = \text{in}_m(I)A$.*

Proof. Theorem 3.6 of [BGS93] proves the statement in the case where $<$ is a monomial order. The same proof also goes through in our context, provided that we take care of the high powers of the negative variables in the same way as we did in the proof of proposition 15. $\square$

We now define the monic functor $\mathcal{H}^{\text{mon}(<,\Delta)}$. Corollary 13 implies that monic ideals are bounded in most cases. We shall later see that ideals arising from the Gröbner functor are always bounded (see lemma 23). It is thus natural, and not very restrictive, to add this boundedness condition when we define functors of monic families.

Definition 18. Let $<$ be a quasi-homogeneous total order and Δ a standard set. Let B be a noetherian k -algebra and $\mathcal{H}^{\text{mon}(<,\Delta)}(B)$ be the set of ideals $I \subset B[\mathbf{x}]$ such that

- I is bounded and Δ -monic
- $B[\mathbf{x}]/I$ is B locally free of rank $\#\Delta$

This defines a covariant functor $\mathcal{H}^{\text{mon}(<,\Delta)}$, which we call a *monic functor*, from the category of noetherian k -algebras to the category of sets.

5. MONIC FUNCTORS ARE REPRESENTABLE

The goal of this section is to prove that monic functors are representable (see theorem 21)

Proposition 19. *Let $<$ be a quasi-homogeneous total order. Let $I \subset B[\mathbf{x}]$ be a bounded ideal with $B[\mathbf{x}]/I$ a locally free B -module of rank n , and let $\mathcal{I} \subset \mathcal{O}_X[\mathbf{x}]$ be the ideal sheaf on $X := \text{Spec } B$ defined by I . Let Δ be a standard set of cardinality n . There exists a locally closed subscheme $Z \subset X$ such that*

- *the restriction of $\mathcal{I}$ to Z is a bounded Δ -monic family,*
- *any morphism $f : \text{Spec } A \rightarrow \text{Spec } B$ such that $IA[\mathbf{x}]$ is a bounded Δ -monic family factors through Z .*

The last proposition is a particular case of the following statement, which we shall prove by induction.

Proposition 20. *Let $<$ be a quasi-homogeneous total order. Let $I \subset B[\mathbf{x}]$ be an ideal bounded by $r_1, \dots, r_d$ such that $B[\mathbf{x}]/I$ is a locally free B -module of rank n , and let $\mathcal{I} \subset \mathcal{O}_X[\mathbf{x}]$ be the ideal sheaf on $X := \text{Spec } B$ defined by I . Let $C := \{x^e : e_i < r_i\}$ be the “hypercube of monomials” of edge lengths r_i , and therefore, of cardinality $s := \prod_{i=1}^d r_i$. We number the monomials $m_i \in C$ such that $m_1 > m_2 > \dots > m_s$. Let $r \leq s$, and fix a map $\mu : \{1, \dots, r\} \rightarrow \{0, 1\}$. Then there exists a locally closed subscheme $Z_r \subset X$ (possibly empty) such that*

- *The sheaf of ideals $\mathcal{I}_r$, which we define as the restriction of $\mathcal{I}$ to Z_r , is a bounded monic family with $\text{in}(\mathcal{I}_r)_m = \langle 1 \rangle$ for $m \notin C$ and $\text{in}(\mathcal{I}_r)_{m_i} = \langle \mu(i) \rangle$ for $1 \leq i \leq r$.*
- *Let $f : \text{Spec } A \rightarrow \text{Spec } B$ be a morphism and $K := IA[\mathbf{x}]$. Then $\text{in}(K)_m = \langle 1 \rangle$ for $m \notin C$ and $\text{in}(K)_{m_i} = \langle \mu(i) \rangle$ for $1 \leq i \leq r$ if, and only if, f factors through Z_r .*

Proof. We start with the first item, which we prove by induction on $r \geq 0$.

When $r = 0$, we may take $Z_0 = X$ since the family is bounded by $r_1, \dots, r_d$, which implies that $\text{in}(\mathcal{I}_r)_{x_i^{r_i}} = \langle 1 \rangle$, hence $\text{in}(\mathcal{I}_r)_m = \langle 1 \rangle$ for $m \notin C$. We may assume that Z_{r-1} is adequately defined. Let $F_r \subset Z_{r-1}$ the closed subscheme defined by the sheaf of ideals $\mathcal{I}(F_r) := (\mathcal{I}_{r-1})_{m_r}$. Let $O_r := Z_{r-1} \setminus F_r$. We define

$$Z_r := \begin{cases} F_r & \text{if } \mu(r) = 0 \\ O_r & \text{otherwise.} \end{cases}$$

By proposition 16 and the induction hypothesis, we have $\text{in}(\mathcal{I}_r)_m = \langle 1 \rangle$ for $m \notin C$ and $\text{in}(\mathcal{I}_r)_{m_i} = \langle \mu(i) \rangle$ for $1 \leq i \leq r-1$. We have to prove that $\text{in}(\mathcal{I}_r)_{m_r} = \langle \mu(r) \rangle$. This is a local problem, so we may assume that $Z_{r-1} \subset X$ is a closed subscheme defined by an ideal $J_{r-1} \subset B$. If $\mu(r) = 1$, the base change $Z_r \hookrightarrow Z_{r-1}$ is open, thus flat. Base change therefore shows that $\text{in}(\mathcal{I}_r)_{m_r} = \langle 1 \rangle$ on a neighborhood of any $p \in Z_r$.

Assume now that $\mu(r) = 0$. We claim that $\text{in}(\mathcal{I}_r)_{m_r} = 0$. The problem is local, so we may assume that both Z_r and Z_{r-1} are affine. Accordingly, we replace the sheaves $\mathcal{I}_r$ and $\mathcal{I}_{r-1}$ by their respective ideals of global sections, which we denote by I_r and I_{r-1} , resp. We will argue by contradiction, supposing that there exists some $f \in I_r$ with $\text{in}(f) = cm_r$, $c \neq 0$. Take some $g \in I_{r-1}$ that restricts to f over Z_r . Since I is bounded by $r_1, \dots, r_d$, we may assume that both f and g are linear combinations of monomials in C , $f = \sum_{m_i \in C} a_i m_i$ and $g = \sum_{m_i \in C} b_i m_i$, resp. Among all possible g , choose one which minimizes $\text{in}(g)$. Then $\text{in}(g) = dm$ with $m \geq m_r$. Suppose that $m > m_r$. Since $\text{in}(I_{r-1})_m = \langle 0 \rangle$ or $\langle 1 \rangle$ by induction hypothesis and since $d \neq 0$, we obtain that $\text{in}(I_{r-1})_m = \langle 1 \rangle$. Choose $h \in I_{r-1}$ with $\text{in}(h) = m$. Then $g' := g - dh$ contradicts the minimality of g . Thus $m = m_r$, and $d \in (I_{r-1})_{m_r} = I(F_r)$ vanishes on F_r . Since d restricts to c on F_r , it follows that $c = 0$. This is a contradiction, which finishes the proof of $\text{in}(\mathcal{I}_r)_{m_r} = 0$.

We now come to the second point. If f factors through Z_r , then $\text{in}(K)_m = \langle 1 \rangle$ for $m \notin C$ and $\text{in}(K)_{m_i} = \langle \mu(i) \rangle$ for $1 \leq i \leq r$, as these properties are inherited from Z_r by proposition 16.

If, on the other hand, f does not factor through Z_r , then we want to prove that $\text{in}(K)_{m_i} \neq \langle \mu(i) \rangle$ for some i , $1 \leq i \leq r$. We may assume that f factors through Z_{r-1} , since in the complementary case, we are done by induction.

We first consider the case $\mu(r) = 0$. By the factorization property of f through Z_{r-1} , we may assume that B is the coordinate ring of the scheme Z_{r-1} . We denote by $f^\# : B \rightarrow A$ the morphism associated with f . Since f does not factor through Z_r , there exists some $g \in I_{r-1}$ with $\text{in}(g) = cm_r$, $f^\#(c) \neq 0$. The pullback of g to $A[\mathbf{x}]$ shows that $\text{in}(IA[\mathbf{x}])_{m_r} \neq \langle 0 \rangle$, and we are done.

Now consider the case $\mu(r) = 1$. Since $Z_r \subset Z_{r-1}$ is open in this case, the factorization property of f implies the existence of a point $p \in \text{Spec } A$ with $f(p) \in Z_{r-1}$ and $f(p) \notin Z_r = Z_{r-1} \setminus F_r$. In particular, $\text{in}(I \cdot k(p)[\mathbf{x}])_{m_r} = 0$ by proposition 16. Thus $\text{in}(K)_{m_r} \neq \langle 1 \rangle$, as expected, since otherwise we would have $\text{in}(I \cdot k(p)[\mathbf{x}])_{m_r} = \langle 1 \rangle$ by proposition 16. $\square$

Theorem 21. *Let $<$ be a quasi-homogeneous total order. Then the monic functor $\mathcal{H}^{\text{mon}(<, \Delta)}$ is representable by a locally closed scheme $H^{\text{mon}(<, \Delta)}$ of $H^n(\mathbb{A}^d)$, where $n = \#\Delta$.*

Proof. Let $I \subset B[\mathbf{x}]$ be an ideal defining a flat family $\text{Spec } B[\mathbf{x}]/I \rightarrow \text{Spec } B$ of relative length n with I bounded and Δ -monic. Let $L_i \subset H^n(\mathbb{A}^d)$ be the closed subscheme of H^n parametrizing the subschemes Z included in the subscheme $\{x_i^n = 0\} \subset \mathbb{A}^d$. Let $L := \bigcap_{x_i \text{ negative}} L_i$. Since the universal ideal of the Hilbert scheme (see [Led11] for the construction and properties of that universal ideal) is bounded over L , there is a locally closed subscheme $L_\Delta \subset L$ parametrizing Δ -monic ideals (see proposition 19). By universal property of the Hilbert scheme, the ideal I corresponds to a unique morphism $\phi : \text{Spec } B \rightarrow H^n(\mathbb{A}^d)$. Lemma 14 implies that the morphism ϕ factors through L . Proposition 19 (the universal property of L_Δ) implies that ϕ even factors through L_Δ . Conversely, any morphism $\text{Spec } B \rightarrow L_\Delta$

yields a Δ -monic bounded ideal by pullback of the universal ideal over the Hilbert scheme. Upon defining $H^{\text{mon}(<, \Delta)} := L_\Delta$, we thus get the required result. $\square$

Proposition 22. *Let $<_j$ be a sequence of quasi-homogeneous total orders converging to $<$. Then for j large, the functors $\mathcal{H}^{\text{mon}(<_j, \Delta)}$ and $\mathcal{H}^{\text{mon}(<, \Delta)}$ are isomorphic.*

Proof. It suffices to prove that the representants $H^{\text{mon}(<_j, \Delta)}$ and $H^{\text{mon}(<, \Delta)}$ are isomorphic. Recall the subscheme L introduced in the proof of the theorem. The subschemes $H^{\text{mon}(<_j, \Delta)}$ and $H^{\text{mon}(<, \Delta)}$ of L are constructed locally using proposition 20. By proposition 11, both constructions coincide locally for $j > j_O$ on an open set O . Since L is quasi-compact, $H^{\text{mon}(<_j, \Delta)} \simeq H^{\text{mon}(<, \Delta)}$ for $j > \max_O j_O$. $\square$

6. BIALYNICKI-BIRULA FUNCTORS ARE REPRESENTABLE

We have proved that our monic functors are representable by schemes. The goal of the present section is to prove that the Białynicki-Birula functors are representable by realizing them as an intersection of two monic functors.

Since monic functors are bounded, we first need:

Lemma 23. *Each ideal $I \in \mathcal{H}^{BB(\xi, \Delta)}(B)$ is bounded with respect to any total order $<_{\xi}$ refining $<_{\xi}$.*

Proof. Up to reordering the components of ξ , one can assume that $\xi_i > \xi_{i+1}$. We denote by k, l the integers such that $\xi_i > 0 \Leftrightarrow i < k$ and $\xi_i < 0 \Leftrightarrow i \geq l$.

By the definition of boundedness, we need to exhibit a family of polynomials h_i . If x_i is a positive variable, we construct h_i as in the beginning of the proof of 10.

Note that the monomials with exponents in Δ form a basis of $B[\mathbf{x}]/I$. In particular, every monomial $\mathbf{x}^e$ leads to an element $\mathbf{x}^e + \sum_{m \in \Delta} c_m \mathbf{x}^m$ of I . For $i \geq l$, $r_i \gg 0$ and $\mathbf{x}^e = x_i^{r_i}$, we have $c_m = 0$ since I is Δ -monic. Thus $h_i := x_i^{r_i} \in I$.

Let x_i be a negative variable of weight 0. To prove that $x_i^r \in I$ for large r , we shall prove that for any monomial $m \in B[x_l, \dots, x_d]$, $x_i^r m \in I$ for large r , and subsequently apply that to the monomial $m = 1$. If the exponent $(e_l, \dots, e_d)$ of $m = x_l^{e_l} \dots x_d^{e_d}$ is large (in more concrete terms, if $e_i \geq r_i$ for some i), then $m \in I$ by the above. We are therefore left with a finite collection of m for which the claim has to be checked. We proceed by induction over m . Let be the m minimal element of the finite collection for which the claim has not been proved yet. By our hypothesis on I , for large r , the monomial $x_i^r m$ lies in the limit ideal $I(0)$, thus I contains an element $f = x_i^r m + R$, where all terms of R are strictly smaller with respect to $<_{\xi}$ than $x_i^r m$. We write the monomials appearing in R as $m_1 m_2 m_3$, with $m_1 \in k[x_1, \dots, x_{k-1}]$, $m_2 \in k[x_k, \dots, x_{l-1}]$, $m_3 \in k[x_l, \dots, x_d]$. Such a product satisfies $m_1 m_2 m_3 <_{\xi} x_i^r m$ only if $m_3 < m$. By induction we know that when multiplying by an adequate power x_i^p , we get $x_i^p m_3 \in I$, hence $x_i^{r+p} m = x_i^p f - x_i^p R \in I$, as required. $\square$

Proposition 24. *Let $\xi \in \mathbb{Z}^d$. Let δ_j be a sequence in $\mathbb{R}^d$ converging to 0 such that the sequence $\xi_j^+ := \xi + \delta_j$ converging to ξ has the property that $<_{\xi_j^+}$ converges to a refinement $<_+$ of $<_\xi$. Let $\xi_j^- := \xi - \delta_j$, and let $<_-$ the limit of $<_{\xi_j^-}$. Then $\mathcal{H}^{BB(\Delta, \xi)} = \mathcal{H}^{\text{mon}(<_+, \Delta)} \cap \mathcal{H}^{\text{mon}(<_-, \Delta)}$.*

Proof. It is clear that $\mathcal{H}^{BB(\Delta, \xi)} \subset \mathcal{H}^{\text{mon}(<_+, \Delta)} \cap \mathcal{H}^{\text{mon}(<_-, \Delta)}$ since by proposition 8, $\mathcal{H}^{BB(\Delta, \xi)} \subset \mathcal{H}^{\text{mon}(<, \Delta)}$ for any refinement $<$ of $<_\xi$. Conversely, take $I \in \mathcal{H}^{\text{mon}(<_+, \Delta)}(B) \cap \mathcal{H}^{\text{mon}(<_-, \Delta)}(B)$. For proving that $I \in \mathcal{H}^{BB(\Delta, \xi)}(B)$, we first prove that $\text{in}_{<_{\xi, m}}(I) = \langle 1 \rangle$ for $m \notin \Delta$.

We argue by contradiction. Suppose that the set $C := \{m \notin \Delta, \text{in}_{<_{\xi, m}}(I) \neq \langle 1 \rangle\}$ is non empty. Since I is bounded, C is finite. Let m' be the smallest element of C for the order $<_+$. Since $m' \notin \Delta$, there is an $f \in I$ with $\text{in}_{<_+}(f) = m'$. We write $f = m' + r + s + t$, with $r = \sum_{f_\xi(e) < f_\xi(m')} c_e \mathbf{x}^e$, $s = \sum_{e \in \Delta, f_\xi(e) = f_\xi(m'), e <_+ m'} c_e \mathbf{x}^e$, $t = \sum_{e \notin \Delta, f_\xi(e) = f_\xi(m'), e <_+ m'} c_e \mathbf{x}^e$. By minimality of m' , for any term $c_e \mathbf{x}^e$ in t , there exists some $g_e \in I$ of shape $g_e = \mathbf{x}^e + \sum_{f_\xi(f) < f_\xi(e)} d_f \mathbf{x}^f$.

Let

$$h := f - \sum_{c_e \mathbf{x}^e \text{ a term of } t} c_e g_e$$

Then h reads $h = m' + r + s + u$ with $u = \sum_{e \notin \Delta, f_\xi(e) = f_\xi(m'), e <_+ m'} c_e (\mathbf{x}^e - g_e)$. The only terms $c_e \mathbf{x}^e$ in h with $f_\xi(e) = f_\xi(m')$ are the terms in s . Note that the orders $<_+$ and $<_-$ have been chosen such that any pair of monomials m, m' satisfies the following:

- if $f_\xi(m) \neq f_\xi(m')$, then $m <_+ m' \Leftrightarrow m <_- m'$
- if $f_\xi(m) = f_\xi(m')$, then $m <_+ m' \Leftrightarrow m' <_- m$.

Thus, if $s \neq 0$, then $\text{in}_{<_-}(h)$ is a term $c_e \mathbf{x}^e$ with $e \in \Delta$, contradicting the assumption that $I \in \mathcal{H}^{\text{mon}(<_-, \Delta)}(B)$. We thus obtain that $s = 0$ and $\text{in}_{<_{\xi}}(h) = m'$, contradicting the definition of C .

Since I is bounded, we get by definition a collection of polynomials $h_i \in I \cap B[x_i]$ with $\text{in}(h_i) = x_i^{r_i}$ when x_i is positive and $h_i = x_i^{r_i}$ if x_i is negative. Then $I(t)$ contains the elements $t^{-\xi_i r_i}(t.h_i) = x_i^{r_i} + \sum_{r < r_i} \lambda_r(t) x_i^r$. It follows that the quotient $B[t, \mathbf{x}]/I(t)$ is a finite $B[t]$ -module.

Summing things up, $\text{Spec } B[t, \mathbf{x}]/I(t)$ is a finite family over $\text{Spec } B[t]$. By construction, this is a flat family of relative length n over the open set $t \neq 0$. The fiber $\text{Spec } B[\mathbf{x}]/I(0)$ over $t = 0$ is a quotient of $\text{Spec } B[\mathbf{x}]/I^\Delta$ which is a flat family of relative length n over $\text{Spec } B$. It follows by semi-continuity that $B[t, \mathbf{x}]/I(t)$ is a locally free $B[t]$ -module of rank n with $I(0) = I^\Delta$. $\square$

Theorem 25. *The Bialynicki-Birula functor $\mathcal{H}^{BB(\Delta, \xi)}$ is representable.*

Proof. By the above, the Bialynicki-Birula functor is an intersection of two functors $\mathcal{H}^{\text{mon}(<_+, \Delta)}$ and $\mathcal{H}^{\text{mon}(<_-, \Delta)}$, both representable by locally closed subschemes

$H^{\text{mon}(<+, \Delta)}$ and $H^{\text{mon}(<-, \Delta)}$, respectively, of the Hilbert scheme. The Białynicki-Birula functor is therefore representable by the schematic intersection $H^{\text{mon}(<+, \Delta)} \cap H^{\text{mon}(<-, \Delta)}$. $\square$

Here is an application of the above to a specific choice of weight ξ .

Proposition 26. *Let $\epsilon : \{1, \dots, d\} \rightarrow \{1, -1\}$ be a sign function. Let ξ be a weight such that $|\xi_1| \gg |\xi_2| \gg \dots \gg |\xi_d| > 0$ and $\text{sign}(\xi_i) = \epsilon(i)$. Let $<$ be the signed lexicographic order defined by $(a_1, \dots, a_d) < (b_1, \dots, b_d)$ if, and only if, $(\epsilon(1)a_1, \dots, \epsilon(d)a_d) < (\epsilon(1)b_1, \dots, \epsilon(d)b_d)$ for the usual lexicographic order. Then $H^{\text{mon}(<, \Delta)}$ is isomorphic to $H^{BB(\xi, \Delta)}$.*

The particular weights of the proposition above are used [ES87], [ES88], [Gro96], [Nak99], for studying the Hilbert scheme of points in the two-dimensional case. But more generally, we have:

Proposition 27. *Let ξ_j be a sequence in $\mathbb{Q}^d$ such that $<_{\xi_j}$ converges to the total quasi-homogeneous order $<$. Then for large j , $\mathcal{H}^{BB(\xi_j, \Delta)}$ is isomorphic to $\mathcal{H}^{\text{mon}(<, \Delta)}$.*

Proof. Following proposition 24, for each ξ_j , we choose a sequence δ_{jk} such that the order $<_{\xi_j + \delta_{jk}}$ (resp. $<_{\xi_j - \delta_{jk}}$) converges to a refinement $<_{j+}$ (resp. $<_{j-}$) of $<_{\xi_j}$, and

$$\mathcal{H}^{BB(\xi_j, \Delta)} = \mathcal{H}^{\text{mon}(<_{j+}, \Delta)} \cap \mathcal{H}^{\text{mon}(<_{j-}, \Delta)}.$$

By proposition, 22, the isomorphisms

$$\mathcal{H}^{\text{mon}(<_{j\pm}, \Delta)} \simeq \mathcal{H}^{\text{mon}(<_{\xi_j \pm \delta_{jk_j}}, \Delta)}$$

hold for k_j large. If we choose k_j large enough, δ_{jk_j} is arbitrarily small compared to ξ_j and we have the convergence of the orders $\lim_j <_{\xi_j \pm \delta_{jk_j}} = \lim_j <_{\xi_j} = <$. In particular, for j large,

$$\mathcal{H}^{\text{mon}(<_{\xi_j \pm \delta_{jk_j}}, \Delta)} \simeq \mathcal{H}^{\text{mon}(<, \Delta)}.$$

The result follows from the displayed equalities. $\square$

7. BIAŁYnicki-BIRULA SCHEMES AND HILBERT-CHOW MORPHISMS

The goal of this section is to prove the following theorem:

Theorem 28. *If $\xi_i \leq 0$, then $H^{BB(\xi, \Delta)}$ is schematically included in the fiber over the origin $\rho_i^{-1}(O)$, where $\rho_i : H^n(\mathbb{A}^d) \rightarrow \text{Sym}^n(\mathbb{A}^1)$ is the Hilbert-Chow morphism associated with the i^{th} coordinate.*

To be precise, let $\rho : H^n(\mathbb{A}^d) \rightarrow \text{Sym}^n(\mathbb{A}^d)$ be the usual Hilbert-Chow morphism. The projection $p_i : \mathbb{A}^d \rightarrow \mathbb{A}^1$ to the i^{th} -coordinate induces a morphism $p_i^n : \text{Sym}^n(\mathbb{A}^d) \rightarrow \text{Sym}^n(\mathbb{A}^1)$. We denote by $\rho_i := p_i^n \circ \rho$ the Hilbert-Chow morphism associated with the i^{th} coordinate. We denote by $O \in \text{Sym}^n(\mathbb{A}^1)$ the point corresponding to n copies of the origin of $\mathbb{A}^1$.

Lemma 29. *Let $I \in \mathcal{H}^{\text{mon}(<, \Delta)}(B)$. Suppose that x_i is a negative variable. Let m_i be the multiplication by x_i in $B[\mathbf{x}]/I$. Then there exists a basis of $B[\mathbf{x}]/I$ such that the matrix of m_i is strictly lower triangular.*

Proof. The monomials b_i with exponents in Δ are a basis of $k[\mathbf{x}]/I$. We order them such that $b_1 > b_2 > \dots > b_n$. Since x_i is negative, whenever $x_i b_j \in \Delta$, then $x_i b_j = b_l$ for some $l > j$. If $x_i b_j \notin \Delta$, we may choose $f \in I$ with $\text{in}(f) = x_i b_j$ and $f = x_i b_j + \sum_{e \in \Delta, \mathbf{x}^e < x_i b_j} c_e \mathbf{x}^e$. Then in the quotient $B[\mathbf{x}]/I$, the identity $x_i b_j = x_i b_j - f = -\sum_{e \in \Delta, \mathbf{x}^e < x_i b_j} c_e \mathbf{x}^e$ holds. $\square$

Proof of the theorem. According to proposition 24, it suffices to prove the inclusion $H^{\text{mon}(<, \Delta)} \subset \rho_i^{-1}(O)$ if the variable x_i is negative for the order $<$.

We recall the observation by Bertin [Ber10] that the Hilbert-Chow morphism is given by the linearized determinant of Iversen. Let $I \in \mathcal{H}^{\text{mon}(<, \Delta)}(B)$ and $b_1, \dots, b_n$ a basis of $B[\mathbf{x}]/I$. If $P \in k[x_i]$, we denote by C_P^j the j^{th} column of the matrix (with respect to our fixed basis) of multiplication by P . If $P_1 \otimes \dots \otimes P_n$ is a pure tensor in $k[x_i]^{\otimes n}$, we put $\text{ld}(P_1 \otimes \dots \otimes P_n) := \det(C_{P_1}^1, \dots, C_{P_n}^n)$. The symmetric group S_n acts on $k[x_i]^{\otimes n}$; we denote by $k[x_i]^{(n)} \subset k[x_i]^{\otimes n}$ the invariant part. Iversen [Ive70] proved that $\text{ld} : k[x_i]^{(n)} \rightarrow B$ is a k -algebra homomorphism. As was remarked by Bertin, this homomorphism corresponds to the Hilbert-Chow morphism ρ_i . The ideal of the origin is generated by the elementary symmetric polynomials, which have degree at least one. For proving the theorem, it therefore suffices to show that $\det(C_{P_1}^1, \dots, C_{P_n}^n) = 0$ if x_i divides some P_j . However, according to the lemma above, that determinant is the determinant of a lower triangular matrix, and that triangular matrix has a zero term on the diagonal if x_i divides some P_j . $\square$

REFERENCES

- [Ber10] José Bertin, *The punctual Hilbert scheme: an introduction*, Séminaires et Congrès **24** (2010), 1–100.
- [BGS93] Dave Bayer, André Galligo, and Mike Stillman, *Gröbner bases and extension of scalars*, Computational algebraic geometry and commutative algebra (Cortona, 1991), Sympos. Math., XXXIV, Cambridge Univ. Press, Cambridge, 1993, pp. 198–215. MR MR1253992 (94j:13021)
- [ES87] Geir Ellingsrud and Stein Arild Strømme, *On the homology of the Hilbert scheme of points in the plane*, Invent. Math. **87** (1987), no. 2, 343–352. MR 870732 (88c:14008)
- [ES88] ———, *On a cell decomposition of the Hilbert scheme of points in the plane*, Invent. Math. **91** (1988), no. 2, 365–370. MR 922805 (89f:14007)
- [Eva04] Laurent Evain, *Irreducible components of the equivariant punctual Hilbert schemes*, Adv. Math. **185** (2004), no. 2, 328–346. MR MR2060472 (2005g:14009)
- [Gro96] I. Grojnowski, *Instantons and affine algebras. I. The Hilbert scheme and vertex operators*, Math. Res. Lett. **3** (1996), no. 2, 275–291. MR 1386846 (97f:14041)
- [Ive70] Birger Iversen, *Linear determinants with applications to the Picard scheme of a family of algebraic curves*, Lecture Notes in Mathematics, Vol. 174, Springer-Verlag, Berlin, 1970. MR 0292835 (45 #1917)
- [Led11] Mathias Lederer, *Gröbner strata in the Hilbert scheme of points*, J. Commut. Algebra **3** (2011), no. 3, 349–404. MR 2826478
- [MS10] Diane Maclagan and Gregory G. Smith, *Smooth and irreducible multigraded Hilbert schemes*, Adv. Math. **223** (2010), no. 5, 1608–1631. MR 2592504 (2011e:14009)

- [Nak99] Hiraku Nakajima, *Lectures on Hilbert schemes of points on surfaces*, University Lecture Series, vol. 18, American Mathematical Society, Providence, RI, 1999. MR 1711344 (2001b:14007)

E-mail address: laurent.evain@univ-angers.fr

UNIVERSITÉ D'ANGERS, FACULTÉ DES SCIENCES, DÉPARTEMENT DE MATHÉMATIQUES, 2, BOULEVARD LAVOISIER, 49045 ANGERS CEDEX 01, FRANCE

E-mail address: mlederer@math.cornell.edu

DEPARTMENT OF MATHEMATICS, CORNELL UNIVERSITY, MALOTT HALL, ITHACA, NEW YORK 14853, USA